

# effects of omega-3 PUFA (ALA) on WT1 gene expression in pancreatic cancer cell line (MIA PaCa-2)

Babak Rahmani (1)  
Dariush Hamed Asl (1)  
Taghi Naserpour Farivar (2)  
Mehdi Azad (3)  
Mehdi Sahmani (4)  
Nematollah Gheibi (2)

1-Department of Molecular Medicine, Faculty of Medical Sciences, Qazvin University of Medical Sciences, Qazvin, Iran

2-Cellular and Molecular Research Centre, Qazvin University of Medical Sciences, Qazvin, Iran

3-Department of Medical laboratory sciences, Faculty of Allied Medicine, Qazvin University of Medical Sciences, Qazvin, Iran

4-Department of Clinical Biochemistry and Genetic, Faculty of Medicine, Qazvin University of Medical Sciences, Qazvin, Iran

Corresponding authors:

Nematollah Gheibi, PhD

Cellular and Molecular Research Center, Qazvin University of Medical Sciences, Qazvin, Iran

Phone: +982833375160

Email: [ngheibi@qums.ac.ir](mailto:ngheibi@qums.ac.ir) ; [gheibi\\_n@yahoo.com](mailto:gheibi_n@yahoo.com)

Mehdi Sahmani, PhD

Mailing address: Dr. M. Sahmani; Department of Clinical Biochemistry and Genetic, Faculty of Medicine, Qazvin University of Medical Sciences, Qazvin – Iran

Phone: +982833328213

Email: [m.sahmani@gmail.com](mailto:m.sahmani@gmail.com)

## Abstract

**Background:** Cancer starts by cells acquiring a number of characteristic alterations especially abnormal cell growth. These properties reflect faults in signalling pathways in cancer cells. Wilms' tumor 1(WT1) acts as a tumor suppressor by negative regulation of WNT/beta-catenin signaling pathway. The role of WT1 in many cancers has been studied. **Aims:** To evaluate the WT1 gene expression alterations in response to different concentrations of omega-3 PUFA alpha linolenic acid (ALA) in pancreatic cancer cell line (MIA PaCa-2) in a time dependent manner. **Methods:** MiaPaca-2 cell line was cultured in monolayers. After that began the cells treatment with omega-3 fatty acid (ALA) using different concentrations of 25, 50, 100, 250, 500 and 1000  $\mu$ M for 24, 48 and 72 hours. The RNA was extracted from both control (untreated) and treated cells. Viability was checked by MTT assay and WT1 expression was evaluated by RT-PCR.

**Results:** Real-time RT-PCR analysis showed that the level of WT1 mRNA, was decreased after 24, 48, 72 hours treatment with omega-3 fatty acid (ALA) for 100, 250, 500 and 1000  $\mu$ M

concentrations but not 25 and 50  $\mu$ M. Data obtained from MTT revealed an-tiliproliferative effects of omega-3 fatty acid (ALA) for 100-1000  $\mu$ M concentrations but not 25 and 50  $\mu$ M.

Conclusion: Our findings indicated that the cell vi-ability and level of WT1 mRNA was decreased dur-ing Omega 3 ALA Treatment. So, it a potential role of Omega 3 in prevention or as a supplementary for pancreatic cancer treatment has been suggested

.Key words: omega-3 PUFA (ALA), pancreatic can-cer, MIA PaCa-2, WT1.

Please cite this article as: Herati E. et al. The effects of omega-3 PUFA (ALA) on WT1 gene expression in pancreatic cancer cell line (MIA PaCa-2).

2018;16(2):275-281. World Family Medicine.

DOI: 10.5742/MEWFM.2018.93287